

Sea Turtle Handling Guidelines

STEP 1:

Determine if the turtle is small enough to bring aboard.

Remember to use gaffs only on fishing gear, not on turtles





IF TURTLE IS TOO BIG TO BRING ABOARD:

- Bring turtle close to boat by pulling gently on the line.
- Determine if turtle is hooked or entangled, and choose the proper tools to remove as much fishing gear as possible from the turtle – including the hook.
- If turtle is hooked and the hook is visible just inside the mouth or on the body, use long handled dehooker to remove hook.
 See Step 3 for instructions.
- If turtle is entangled or the hook is deep inside mouth or throat and cannot be removed, use a long-handled line cutter to cut all lines.
- Skip to Step 5.

IF TURTLE IS SMALL ENOUGH TO BRING ABOARD:

- Use dip net to bring turtle aboard.
- Do not bring turtle aboard by pulling on fishing line or by grabbing the eye sockets.
- It may be helpful to grab the front flippers close to the turtle's body when using the net to help bring it aboard.
- Go to **Step 2**.






STEP 2:

After the turtle has been brought aboard, determine if it is alive or appears dead.

A turtle that looks dead may just be very tired, and can regain strength with your help.



UNCONSCIOUS TURTLE - inactive and appears dead

- Keep the turtle on a tire in a secure, shaded place away from activity.
- · Remove fishing gear using instructions in Step 3.
- Place turtle on its belly and elevate back flippers at least 6 inches for at least 4 hours to help remove water from its lungs while recovering.
- Place a wet towel on turtle. Do not cover nostrils. Occasionally wet turtle with a deck hose. Avoid spraying turtle's head.
- Perform a reflex test every 3 hours, by gently touching corner of eye and lightly pulling on tail. Movement may indicate the turtle is recovering.
- If there is no movement from reflex tests after at least 4 hours, but no more than 24 hours, release the turtle to the ocean using methods in **Step 4**.

CONSCIOUS TURTLE - active or awake

- Keep the turtle on a tire in a secure, shaded place away from activity.
- Remove fishing gear using instructions in Step 3.
- Release the turtle using methods in **Step 4**. You do not have to wait 4 or more hours before release.

Sea Turtle Handling Guidelines

When to Leave Hooks in Place.

It is normally best to remove all fishing gear from the turtle, but there are situations when the gear should not be removed. Leave hook in place and cut line as close as possible to hook <u>if</u>:


The hook has been swallowed.

Forcing a dehooking device down a turtle's throat may worsen its injuries or cause an infection.




The hook has penetrated the roof of the mouth.

Trying to remove this hook may cause damage to the organs and



The hook is in the glottis.

The glottis is located at the back of the mouth and covers the airway. Attempting to remove hooks from the glottis may cause further damage and prevent the turtle from covering its airway during dives.



STEP 3:

Methods for removing fishing gear from a hooked turtle.

TURTLE HOOKED WITH BARB EXPOSED

- 1. Using bolt cutters, remove the barb of the hook.
- 2. Once barb has been clipped off, back the hook out to remove it.

TURTLE HOOKED WITH BARB EMBEDDED

- 1. Follow instructions on using a pig-tail dehooker.
- 2. If hook cannot be removed, cut line as close as possible to hook.

TURTLE IS HOOKED, BUT YOU CANNOT SEE HOOK


• Cut fishing line as close as possible to hook without pulling hard on line.

TURTLE ENTANGLED IN FISHING LINE

Use monofilament or wire cutters to remove all fishing line from turtle.

IF BOLT CUTTERS ARE NOT AVAILABLE

- Place a J-style dehooker or similar hand-held tool on the leader to cut hook.
 A short handled pig-tail dehooker can also be used. To get dehooker on the line, refer to pig-tail dehooker instructions.
- 2. Slide device down the leader to the bottom of the hook.
- 3. Pull the line so it is opposite from the handle of the dehooker.
- 4. Keep the line tight, then pull and twist dehooker to remove hook.



Sea Turtle Handling Guidelines

Dehooking a turtle using a pig-tail dehooker with your right hand:



Place the dehooker at 90° to the line with the end of the pig-tail facing up.



Draw the dehooker back towards you like a bow and arrow until loop pulls on line, maintaining contact between the dehooker and the line.



Rotate the dehooker 1/4 turn clockwise. (The line should be inside the curl of dehooker)



Run the dehooker down the line until it engages the bottom bend of the hook.








Pull the line tight and parallel to the dehooker. Give quick thrusts to remove the hook.



Keep line tight, so the hook remains inside the curl of dehooker, until hook is clear of the turtle.

Circle Hooks



When using the long-handled pig-tail dehooker on a turtle in the water, it may be easier to remove circle hooks if the line is not parallel to the dehooker's handle once the dehooker is on the line.

- While keeping the line tight, separate the line and dehooker, then try to push and twist the dehooker to
- dislodge the hook. This may work better than quick thrusts.
- If you cannot remove the hook, cut line as close as possible to the hook.
- If turtle is aboard, try to rotate the hook back out by using the line or pliers before using the dehooker. This may help remove the hook.

STEP 4:

Carefully return turtle to water.

- 1. Stop vessel, and take engine out of gear.
- 2. Release the turtle away from any fishing gear in the water.
- 3. Gently put turtle in the water, head-first.
- 4. Make sure turtle is clear of vessel before motoring away.

STEP 5:

Record the interaction in your logbook.


- 1. Record the identified species.
- 2. Write down how much fishing gear remained on turtle after release.
- 3. Record any tag numbers observed on turtle.



....

LEATHERBACK

- · Dark, leathery skin covers body and shell
- · No scutes or scales like other turtles
- 5-7 head to tail ridges on back
- Adults are much larger than other turtles





Hard Shell Turtles

GREEN

- 1 pair of prefrontal scales
- · 5 central scutes
- 4 pairs of lateral scutes
- 4 inframarginal scutes on each side





HAWKSBILL

- 2 pairs of prefrontal scales
- · 5 central scutes
- · 4 pairs of overlapping lateral scutes
- · 4 inframarginal scutes on each side









LOGGERHEAD

- · 2 pairs of prefrontal scales
- 5 central scutes
- 5 pairs of lateral scutes
- · 3 inframarginal scutes on each side



OLIVE RIDLEY


- · 2 pairs of prefrontal scales
- 5 central scutes
- 5-9 pairs of lateral scutes
- 4 inframarginal scutes on each side with pores




Leatherback Sea Turtle

(Dermochelys coriacea)


- · Only soft-shelled species
- Dark gray or black with variable white spotting
- 5-7 head-to-tail ridges on back
- Leathery shell
- No scales
- · W-shaped upper jaw or beak
- May attain great size







Approximate range



TURTLE FACTS:

Leatherback turtles interact with both the Hawaii swordfish (shallow-set) and tuna (deep-set) longline fisheries. They are usually hooked or entangled externally, rather than in the mouth. This turtle has a firm, leathery skin covering the shell and body, instead of a hard shell and scales like other turtles. They are highly migratory, swimming long distances across the Pacific from nesting to foraging areas. Leatherbacks are the largest of all sea turtles with adults reaching 6½ feet (2 meters) in length and over 1,500 pounds (681 kg).

Loggerhead Sea Turtle


(Caretta caretta)

2 pairs of prefrontal scales


5 central scutes

5 pairs of lateral scutes

3 pairs of inframarginal scutes



Approximate range






TURTLE FACTS:

Loggerhead turtles interact with both the Hawaii swordfish (shallow-set) and tuna (deep-set) longline fisheries. In the North Pacific, juveniles hatched from nests in Japan swim across the ocean to feed and grow near the Mexican coast. They can spend decades in migratory and developmental habitats in Mexico and the central Pacific until maturity, when they return to Japan. Loggerheads can grow to over 36 inches (92 cm) in shell length and 250 pounds (113 kg). They have large heads, strong jaws, and typically have shells that are reddish-brown with a yellow underside.

Olive Ridley Sea Turtle

(Lepidochelys olivacea)


- 2 pairs of prefrontal scales
- 5 central scutes
- 5-9 pairs of lateral scutes
- 4 pairs of inframarginal scutes with one pore on each scute




Approximate range



TOP VIEW OF HEAD





OF TURTLE

TOP VIEW



TURTLE FACTS:

Olive ridley turtles interact primarily with the Hawaii tuna (deep-set) longline fishery, and occasionally with the swordfish (shallow-set) fishery. These turtles are highly migratory and usually live in warm, tropical waters, but may also occur in cooler waters north of Hawaii. Olive ridley turtles are the smallest sea turtles, averaging 25 inches (61 cm) in shell length and 100 pounds (45 kg). Their shell is generally olive green with a light yellow underside. They sometimes have more lateral scutes on one side of their shell than the other. They are the only turtles in the Pacific with a pore on each inframarginal scute.

Green Sea Turtle


(Chelonia mydas)

1 pair of prefrontal scales


5 central scutes

4 pairs of lateral scutes


4 pairs of inframarginal scutes




Approximate range



TOP VIEW OF HEAD



TOP VIEW



BOTTOM VIEW OF TURTLE



TURTLE FACTS:

The green turtle is the most widespread and commonly-known sea turtle in tropical and sub tropical waters. Green turtles are not usually caught in longline fisheries, but interactions can occur in the Hawaii and American Samoa fisheries. Green turtles are the largest of the hard-shell turtle species and can grow up to 47 inches (120 cm) in shell length and weigh over 300 pounds (136 kg). They get their name from the color of their fat. The shell color can range from yellow-green to reddish-brown to almost black. Loggerheads and olive ridley turtles can be easily mistaken for a green turtle. When in doubt, look at the head and check the number of prefrontal scales.


Hawksbill Sea Turtle

(Eretmochelys imbricata)


- 2 pairs of prefrontal scales
- 5 central overlapping scutes
- 4 pairs of overlapping lateral scutes
- 4 pairs of inframarginal scutes

TOP VIEW

OF HEAD



Approximate range



BOTTOM VIEW

OF TURTLE

TOP VIEW



TURTLE FACTS:

There has been no reported interaction between a hawksbill turtle and the Hawaii longline fisheries. Hawksbills can be found in tropical and sub-tropical regions across the Pacific. They nest in low numbers throughout the Pacific, including Hawaii. Adult hawksbills can grow to almost 3 feet (90 cm) in shell length and weigh up to 200 pounds (91 kg). The hawksbill is named for its sharp, pointed, bird-like beak. The shell has "tortoise shell" coloring, ranging from dark to golden brown, with streaks of orange, yellow, brown and black. These are the only sea turtles that have overlapping scutes on the top shell, like roof shingles.

Marine Mammal Handling and Release Guidelines

Have an identification guide and paper available in case of a marine mammal interaction.

Small Whales and Dolphins



- 1. Make sure the crew is ready to help.
- 2. Move the boat carefully, stop the boat, and put the transmission in neutral when the animal is close.
- 3. If the far side of the mainline is within reach, use gaffs to grab only the line. This will keep any remaining gear in the water from pulling on the line and the animal. Do NOT use gaffs or sharp objects to grab an animal.
- 4. Slowly bring the animal next to the boat.
- 5. Avoid sudden actions that may scare the animal.

If the animal is tangled in line:

- 1. Grab the far side of the mainline and tie the mainline to the boat.
- 2. Use a long-handled line cutter to cut as much line off the animal as you can.

If the animal is hooked:

- 1. Use a dehooker to remove the hook.
- 2. If the hook cannot be removed, use a long-handled line cutter to cut the line as close as you can to the hook.
- 3. Remove as much line as possible from the animal. Do not use ropes or other lines to tie the animal to the boat.

Large Whales

If a large whale is alive and hooked or entangled in fishing gear, immediately call the Disentanglement Hotline at **1-888-256-9840** or the U.S. Coast Guard on VHF Ch. 16 for instructions.

If a large whale is dead and hooked or entangled in fishing gear, immediately call the U.S. Coast Guard on VHF Ch. 16 for instructions.

For All Interactions



Write down as much information as possible to describe the animal, and its injuries.

- How long was the animal?
- What did the animal look like (did it have stripes, spots, or different colors)?
- Was there any fishing gear still on the animal when it was released?
 If so, where, what kind, how much?
- Did you see any tags on the animal? If yes, can you see any letters or numbers on the tag?


After an interaction with a marine mammal, get the rest of your fishing gear out of the water. Then record all the information about the interaction on your Marine Mammal Injury / Mortality Reporting Form, even if you had an observer aboard. Mail the form when you get to port.

If you have a marine mammal interaction, you could have another if you keep fishing in the same area. Move away from the area, and call other fishermen to warn them. If you stay in the same area, wait 2 days before setting your gear to avoid more interactions.


Seabird Handling Guidelines

PLEASE NOTE: If bird is a short-tailed albatross, follow special guidelines for handling short-tailed albatross. For all other seabirds, see below.

1. Stop vessel to reduce tension on the line and bring bird aboard using a dip net.



- Working with another person, hold the back of the bird's head and isolate the hooked or entangled area while the other person takes the bird from the net. Fold the bird's wings to their natural resting position against the body.
- 3. Wrap the bird's wings and feet with a clean towel or blanket. Do not wrap the bird's body too tightly or block the nostrils, as these will prevent the bird from breathing.
- 4. Cut and remove all fishing line from bird. If bird is lightly hooked in the bill, leg, or wing, and the barbed end of the hook is visible, use bolt cutters to cut the barb and then back the hook out. If bird has been deeply hooked, cut the line as close as possible to hook and leave hook in place.



Never attempt to remove a hook from anywhere on a bird by pulling on line.

- 5. Allow bird to dry for 1/2 hour to 4 hours in a safe, enclosed place. Refer to **Release Guidelines.**
- 6. Record any leg band numbers observed on the bird in logbook.



Wear gloves, long sleeves and protective eyewear when handling seabirds. They have sharp beaks and give painful bites.

Release Guidelines

A bird is ready for release when its feathers are dry.



Albatross NOT ready to be released



Albatross ready to be released

If bird is ready for release:

Stop the vessel. Gently place bird onto the surface of the water. Do not throw bird in air or motor away if bird is not clear of vessel.

Short-Tailed Albatross Handling Guidelines

Short-tailed albatross are an endangered species and have special handling requirements.

If you catch a short-tailed albatross:

Immediately try to contact National Marine Fisheries Service, U.S. Coast Guard, or U.S. Fish and Wildlife Service. They will contact an expert to give you advice in the handling and release of short-tailed albatross.

National Marine Fisheries Service (NMFS) (808) 944-2200

U.S. Coast Guard (USCG) 08240.0 KHz (Daytime ITU Channel 816) 12242.0 KHz (Daytime ITU Channel 1205) 04134.0 KHz (Nightime ITU Channel 424) 06200.0 KHz (Nightime ITU Channel 601)

U.S. Fish and Wildlife Service at French Frigate Shoals (USFWS) Contact frequency: 10.0054

If a short-tailed albatross is hooked or entangled:

- 1. Stop vessel to reduce tension on the line and bring bird aboard using a dip net.
- 2. Wrap the bird's wings and feet with a clean towel to protect its feathers from oils or damage.
- 3. Remove any entangled lines from the bird and determine if the bird is dead or alive.

If dead, notifiy NMFS. Label the bird, put it in a plastic bag and store in freezer. Give bird to NMFS when you return to port.

If alive, place bird in a safe, enclosed place and immediately contact NMFS, USCG and USFWS.

If unable to make contact for 24-48 hours, determine if the bird is lightly, moderately, or deeply hooked. See description.

- If bird is deeply hooked, keep bird in a safe, enclosed place until further instructed.
 Do NOT release the bird.
- 5. If bird is lightly or moderately hooked, remove hook by cutting the barb and backing hook out.
- 6. Allow bird to dry for 1/2 hour to 4 hours in a safe, enclosed place. Refer to **Release Guidelines.**
- 7. Record information in the short-tailed albatross recovery data form.



Short-tailed albatross fly across the entire North Pacific. Around Hawaii, only young short-tailed albatross (shown above) have been seen. The number of birds is increasing, but fewer than 3,000 birds remain in the wild.

Is the bird lightly, moderately, or deeply hooked?



Lightly Hooked: Hook is clearly visible on bill, leg or wing.

Moderately Hooked: Hooked in the mouth or throat with hook visible.



Deeply Hooked: Hook has been swallowed and is located inside the bird's body below the neck.

Release Guidelines

The bird is ready for release if it meets ALL of the following criteria:

- Stands on both feet with toes pointed forward
- Holds its head erect and responds to sound and motion
- · Breathes without making noise
- Flaps and retracts wings to normal folding position
- Feathers are dry

If any of these conditions are not met, the bird cannot be released.

If bird is ready for release:

Stop the vessel. Gently place bird onto the surface of the water. Do not throw bird in air or motor away if bird is not clear of vessel.

Seabird Identification

Laysan Albatross (Phoebastria immutabilis)

Feathers: · White head, neck, and belly

> Dark brown upper wings and back · Brown and white under wings

· Dark area around each eve

Legs/Feet Color: Pink to gray

Bill Color: Yellow-pink with gray tip



Black-footed Albatross (Phoebastria nigripes)

Feathers: · Dark brown head, body, and wings

> • Small white patch behind the eyes · White ring around base of bill

· Adults - small white patch at base of tail

Legs/Feet Color: Black-brown

Bill Color: Black-brown





Short-tailed Albatross (Phoebastria albatrus)

ENDANGERED SPECIES

JUVENILE

Feathers: Dark brown head, body, and wings

Legs/Feet Color: Brown-gray

Bill Color: Bright pink with a thin black line around base



SUB ADULT

Feathers: White neck, belly, and back

Dark brown cap and back of neck

· Black and white wings

Legs/Feet Color: Pink to gray

Bill Color: Bright pink with a thin black line around base



ADULT

Feathers: Golden-yellow head and neck

· White back, base of tail, and belly

· Black and white wings

Legs/Feet Color: Pink to gray

Bill Color: Bright pink with a thin black line around base