



Odontocete bycatch and depredation in longline fisheries: A review of available literature and of potential solutions

DEREK J. HAMER,¹ SIMON J. CHILDERHOUSE and NICK J. GALES, Australian Marine Mammal Centre, Australian Antarctic Division, 203 Channel Highway, Kingston, Tasmania 7050, Australia.

ABSTRACT

Operational interactions between odontocetes (*i.e.*, toothed whales) and longline gear are a global phenomenon that may threaten the conservation of odontocete populations and the economic viability of longline fisheries. This review attempts to define the issue, summarize the trends and geographical extent of its occurrence over the last half century, explore the potential impact on odontocetes and on fisheries, and describe potential acoustic and physical mitigation solutions.

Reports of odontocete bycatch rates are highly variable (between 0.002 and 0.231 individuals killed per set) and at least 20 species may be involved. Information about marine mammal population size, migration patterns and life history characteristics are scarce, although at least one population may be in decline due to losses attributable to longline bycatch. Information about the financial impact of depredation on pelagic longline fisheries is also scarce, although estimates of daily fleet-wide losses range between US\$1,034 and US\$8,495 (overall fleet income was not reported). Such biological and financial losses may be unsustainable.

Recent developments in acoustic and physical mitigation technologies have yielded mixed results. Acoustic mitigation technologies have no moving parts, although require complex electronics. To date, they are insufficiently developed and their efficacy has been difficult to assess. Physical mitigation technologies generally require complex moving parts, although they are relatively simple to develop and assess. Further development and testing remains necessary before widespread implementation would be possible. Development of these approaches should be prioritized and a “toolbox” of various strategies and solutions should be compiled, because a single panacea to the problem is unlikely to emerge.

Key words: acoustic, bycatch mortality, depredation, fishing, longline, operational interactions, odontocete, physical, toothed whale.

INTRODUCTION TO OPERATIONAL INTERACTIONS BETWEEN ODONTOCETES AND LONGLINES

The occurrence of operational interactions between cetaceans (*e.g.*, whales, dolphins, and porpoises) and commercial fisheries has attracted considerable attention in the published literature and is a familiar problem to many fishing, management, and research communities (*e.g.*, Northridge 1984, 1991; Beverton 1985; Reeves

¹Corresponding author (e-mail: derek.hamer@aad.gov.au).

et al. 1994; Northridge and Hofman 1999; Donoghue *et al.* 2003; Shaughnessy *et al.* 2003; Read 2005; Gilman *et al.* 2006). Operational interactions involve the simultaneous physical convergence of cetaceans and commercial fisheries upon the same spatially retracted area, often when both are in pursuit of the same fish (Northridge and Hofman 1999, Shaughnessy *et al.* 2003, Hamer *et al.* 2008, Moreno *et al.* 2008). Positive outcomes include (1) fisheries using cetaceans to indicate the presence of fish (Gosliner 1999, Northridge and Hofman 1999) and (2) cetaceans using fisheries to access an otherwise inaccessible food resource (Gilman *et al.* 2006, Moreno *et al.* 2008). Negative outcomes include threats to the viability of (1) cetacean populations when depredating individuals become bycatch and are injured or killed (Gosliner 1999, Shaughnessy *et al.* 2003, Hamer *et al.* 2008) and (2) fisheries when depredating whales remove or damage the catch (Hucke-Gaete *et al.* 2004, Ramos-Cartelle and Mejuto 2008). The negatives may be exacerbated by an increase in the level of trophic interactions due to increased competition for the same fish stock, which results in either direct reduction (through removal of fish), or indirect reduction (through trophic cascades) of target fish stocks (Northridge and Hofman 1999, Kaschner 2004). Both scenarios could reduce the overall quantity of fish available to cetacean populations (Kaschner 2004, Bakun *et al.* 2009) and fisheries (Ashford *et al.* 1996, Earle 1996), thus increasing the likelihood of operational interactions.

A growing body of information concerning the nature and extent of operational interactions between odontocetes (*i.e.*, toothed whales, dolphins, and porpoises) and longline gear has been emerging in the literature, since longlining commenced modernization in the 1950s (Yamaguchi 1989, Ward and Hindmarsh 2007). The main areas identified to be of concern are depredation and bycatch. Depredation occurs when an individual odontocete partially or completely consumes caught fish from the longline, or deters free swimming fish that may otherwise have become caught (Yano and Dahlheim 1995, Northridge and Hofman 1999, Read 2005, Gilman *et al.* 2006, Lauriano *et al.* 2009). Although depredation of bait has also been identified as an issue, there is insufficient data and information currently available to warrant in-depth consideration here. Bycatch occurs when a depredating odontocete becomes caught on a longline hook when attempting to remove the catch (Beverton 1985, Shaughnessy *et al.* 2003, Read 2005, Secchi *et al.* 2005, Indian Ocean Tuna Commission 2007).

The modernization of longlining mid-last century resulted in rapid geographic expansion, with reports of odontocetes depredating catch from longlines emerging soon after (*e.g.*, Iwashita *et al.* 1963, Sivasubramaniam 1964, Mitchell 1975). The growing number of reported incidences since that time suggests the phenomenon may have become a significant economic problem for affected longline fisheries and a significant conservation and welfare problem for affected odontocete populations. This review attempts to (1) define the issue, (2) summarize the available literature to determine temporal trends and geographical extent of catch depredation by and bycatch of odontocetes, (3) explore the impacts of depredation on fisheries and bycatch on odontocete populations, and (4) describe the acoustic and physical tools being developed to mitigate the problem.

SUMMARY OF ODONTOCETE DEPREDAATION AND BYCATCH REPORTS IN THE REVIEWED LITERATURE

The compilation and interpretation of the accessible literature may be useful in revealing the nature and extent of the problem and may provide insights into

how to mitigate the impact of one or both. The literature cited was restricted to peer reviewed documents (articles and reports) that referred to fishery logbook or observer data, specifically relating to operational interactions between odontocetes and longline fisheries. Electronic search engines and databases were used, such as Web of Science, Current Contents, Google Scholar, and general internet searches, using keywords such as: whale, cetacean, odontocete (and individual species names), depredation, and bycatch.

The literature search identified 32 peer reviewed documents matching the specified criteria, published between 1964 and 2010 (Table 1). Early documents merely acknowledged the occurrence of catch depredation, with the first specific account of an odontocete being bycaught on a longline hook emerging in 1983 (Di Natale and Mangano 1983). Nonetheless, the literature has remained focused on the effects on the fishery, with 23 reports of depredation compared with 12 reports of bycatch (Table 1). Twenty-two reports have emerged since 2000, amounting to over twice the number produced over the previous four decades combined (Fig. 1). This recent spike suggests an increase in awareness and interest in the issue.

The literature cited indicates that 20 odontocete species have been involved in operational interactions with longline gear. Fifteen species were confirmed to have either depredated from, or have become bycaught on, longline hooks (Table 1). The five remaining species (*i.e.*, rough toothed dolphin, *Steno bredanensis*; spinner dolphin, *Stenella longirostris*; Atlantic humpback dolphin, *Sousa teuszii*; melon-headed whale, *Peponocephala electra*; and pygmy killer whale, *Feresia attenuata*) were mentioned in the literature cited, but were involved in unverified, anecdotal and unquantified events (Northridge 1984, Nishida and Tanio 2001, South Pacific Regional Environment Program 2002, Culik 2004, Secchi *et al.* 2005, Watson and Kersletter 2006, Moore *et al.* 2010). Based on the literature obtained, killer whales (*Orcinus orca*) and sperm whales (*Physeter macrocephalus*) appear to be the main species involved with demersal longline fisheries at higher latitudes, while false killer whales (*Pseudorca crassidens*) and pilot whales (*Globicephala* spp.) appear to be the main species involved with pelagic longline fisheries at lower latitudes. The problem also appears to be geographically widespread, with reports of depredation from and bycatch on longlines confirmed in 25 locations, from the equator to high latitudes in both hemispheres and in all of the world's major oceans (Fig. 2).

Some key events may explain the recent increase in the number of reports emerging in the available literature. Since the 1940s and 1950s, some odontocete populations appear to have benefited from increased international protection instruments, such as the Convention on International Trade in Endangered Species (CITES) and the International Convention for the Regulation of Whaling (ICRW). During the same period, fishing effort has increased to meet the demands of a burgeoning human population (United Nations 2009) with changing dietary needs (Duarte *et al.* 2009). This situation is likely to have increased the probability of odontocetes encountering fishing gear, thus resulting in increased incidences of depredation and bycatch (Northridge 1984, 1991; Jefferson 1994; Reeves *et al.* 1994; South Pacific Regional Environment Program 2002; Donoghue *et al.* 2003; Gilman *et al.* 2006). As such, the recent growth in the volume of literature may reflect an increase in the motivation of fishermen to find ways of mitigating catch depredation, in a bid to improve or maintain catch returns, at a time when increased operational costs (*i.e.*, fuel and freight) and depleted fish stocks (*i.e.*, overfishing) are eroding profits (Northridge and Hofman 1999, Ebert *et al.* 2009, Food and Agriculture Organization 2009). The emergence of this information has encouraged relevant conservation and management organizations to characterize the problem and to explore mitigation strategies

Table 1. Summary of operational interactions (*i.e.*, catch depredation and bycatch) between odontocetes and pelagic and demersal longline fishing gear, inferred from or quantified in the literature reviewed.

Whale species involved	Fishery details			Catch depredation details		Whale bycatch details			Year
	Species targeted	Region of interaction	Gear configuration	Percent of sets affected ^c	Percent of catch damaged ^c	Percent of whales hooked	Rate (animals/set)	Source	
?	?	IO	?		<55			Sivasubramaniam	1964
KW, FKW	T	IO	P					Mitchell	1975
GTB	?	TC	?			2 ^f		Watson	1981
CD	S	FAC	P			2 ^f		Duguay and Husenot	1982
SW		CM	P			1		Di Natale and Mangano	1983
KW	SF	PWS	D		25			Matkin	1986
KW	SF	BS, PWS	D	(15–25)				Dahlheim	1988
KW	SF, GT, AF	BS, GA	D		(13–45)			Yano and Dahlheim	1995
SW, KW	PT	SG	D	93	>90	2 ^f	0.07	Ashford <i>et al.</i>	1996
KW	T, SwF	SB	P		(50–100)			Secchi and Vaske	1998
KW	SS, BET	NZ	D		5–10			Visser	2000
SRW			P			3 ⁸		Best <i>et al.</i>	2001
SW	SF	GA	D		23 ^c			Straley <i>et al.</i>	2002
KW, B	T, SwF	EA	P			2 ^f		Shaughnessy <i>et al.</i>	2003
SW, KW	PT	SC	D	16	3 (0–100)			Hucke-Gaete <i>et al.</i>	2004
SW, KW	PT	SG	D	13				Perves <i>et al.</i>	2004
KW, PW	BET, L	SAus	D	6–80 ^d				AFMA ⁿ	2005

KW, FKW, B, D	T, SwF	EA	P		5 ^h		Bell <i>et al.</i>	2006
SW, KW	PT	SG, PEI	D	>50			Kock <i>et al.</i>	2006
BD	KM	F	P	6–20			Zollert and Read ⁿ	2006
KW	T, SwF	SB	P	12 (1–47)			Dalla Rosa and Secchi	2007
KW	T, SwF, S	SA	D	0.50			Williams <i>et al.</i>	2007
KW	PT	CA	D	42			Roche <i>et al.</i>	2007
Various ^a					67 ⁱ	0.003 ^k	Forney and Kobayashi	2007
FKW	T, SwF, S	B, AA	P	(1–9)	2 ^f		Hernandez-Milian <i>et al.</i>	2008
FKW	SwF	A, IO, P		2	18 ^j	0.002 ^k	Ramos-Cartelle and Mujeto	2008
SW	SF	BS, GA, AI	D	<1			Sigler <i>et al.</i>	2008
SW, KW	PT	SC	D	0.36			Moreno <i>et al.</i>	2008
SW, KW	PT	CA		41			Tixier <i>et al.</i>	2009
CD, BD, SD	Various ^b	IC	D, P	40			Lauriano <i>et al.</i>	2009
FKW, PW	T, BF	CS	P	<16	3 ^f	0.231	Hamer	2009 ^b
DD	D,S	PC	P	<16	1	0.05 ^l	Mangel <i>et al.</i>	2010

Continued

Table 1. (Continued)

Whale species abbreviations:
B, Unidentified baleen whale species (Mysticeti); BBW, Blainville's beaked whale (<i>Mesoplodon densirostris</i>); BD, Bottlenose dolphin (<i>Tursiops truncatus</i>); BW, Bryde's whale (<i>Balaenoptera edeni</i>); CD, Common dolphin (<i>Delphinus delphis</i>); D, Unidentified small toothed whale species (Odontoceti); DD, Dusky dolphin (<i>Lagenorhynchus obscurus</i>); FKW, False killer whale (<i>Pseudorca crassidens</i>); GTB, Ginko-toothed beaked whale (<i>Mesoplodon ginkgodens</i>); HW, Humpback whale (<i>Megaptera novaengliae</i>); KW, Killer whale (<i>Orcinus orca</i>); PW, Pilot whale (<i>Globicephala</i> spp.); RD, Risso's dolphin (<i>Grampus griseus</i>); SD, Striped dolphin (<i>Stenella coeruleoalba</i>); SPD, Pantropical spotted dolphin (<i>Stenella attenuata</i>); SRW, Southern Right Whale (<i>Eubalaena australis</i>); SW, Sperm whale (<i>Physeter macrocephalus</i>).
Fish species abbreviations:
AF, Arrowtooth flounder (<i>Atheresthes stomias</i>); BET, Blue-eye trevallia/bluenose (<i>Hyperoglyphe antarctica</i>); BF, Billfish (Istiophoridae and Xiphiidae); D, Dorado (<i>Coryphaena hippurus</i>); GT, Greenland turbot/halibut (<i>Reinhardtius hippoglossoides</i>); KM, King mackerel (<i>Scomberomorus cavalla</i>); L, Unspecified ling species (<i>Genypterus</i> spp.); PT, Patagonian toothfish (<i>Disostichus eleginoides</i>); S, Unspecified shark species (Selachimorpha); SE, Sablefish (<i>Anoplopoma fimbria</i>); SS, School shark (<i>Galeorhinus galeus</i>); SWF, Swordfish (<i>Xiphias gladius</i>); T, Tuna (<i>Thunnus</i> spp.).
Region abbreviations:
A, Atlantic; AA, Azores Archipelago; AI, Aleutian Islands; B, Brazil; BS, Bering Sea; CA, Crozet Archipelago; CM, central Mediterranean; CS, Coral Sea; EA, eastern Australia; F, Florida; FAC, French Atlantic coast; GA, Gulf of Alaska; IC, Italian coast; IO, Indian Ocean; NZ, New-Zealand; P, Pacific; PC, Peruvian coast; PEI, Prince Edward Island; PWS, Prince William Sound; SA, South Africa; SAus, southern Australia; SB, southern Brazil; SC, southern Chile; SG, South Georgia; TC, Taiwanese coast.
Additional details:
^a PW, FKW, SPD, BD, BBW, RD, SW, BD, CD, and HW.
^b Unspecified fish species.
^c Values are averages or estimates; values in parentheses are ranges.
^d 6% of sets affected, calculated from industry data; 80% of sets affected, derived from anecdotal information from fishermen.
^e Inferred from a reduction in the catch rate of the targeted fish.
^f Dead animals recorded.
^g Entanglement mortalities.
^h 5 animals hooked; 2 dead (1 KW and 1 D) and 3 released alive.
ⁱ 67 hooked; 7 dead (2 PW, 2 FKW, 1 SPD, 1 BD, and BBW) and 60 released alive.
^j 18 animals hooked; proportion dead and released alive not specified.
^k Derived retrospectively from figures presented in the results of the study.
^l In addition, harpooning of dolphins for bait was occasionally observed.
^m Australian Fisheries Management Authority.
ⁿ Study of a troll fishery—included here due to the relevance of the depredation mitigation strategy to longline fishing.




Figure 1. Decadal summary of the number of reviewed reports of operational interactions between odontocetes and longline gear over the last 50 yr (the 2000s includes one study published in early 2010).

that facilitate the continued conservation of recovering cetacean populations and to minimize mounting pressure on the economic viability of fisheries. Mitigation of odontocete bycatch and catch depredation has been prioritized by some fisheries in recent times, indicating that stakeholders rank its importance highly, relative to other issues that impact fishery viability (Donoghue *et al.* 2003, Australian Fisheries Management Authority 2005).

IMPACTS OF BYCATCH ON ODONTOCETES

Longline gear poses a significant injury and drowning risk to depleting odontocetes, which affect the welfare of individuals and the conservation of populations (Ashford *et al.* 1996, Northridge and Hofman 1999, Visser 2000, South Pacific




Figure 2. Geographic distribution of operational interactions (inferred and quantified) between odontocetes and longline gear (gray areas) in accessible reports.

Regional Environment Program 2002, Secchi *et al.* 2005, Gilman *et al.* 2006, Forney and Kobayashi 2007, Hamer 2009a, Lauriano *et al.* 2009, Reeves *et al.* 2009). Some individuals may accidentally ingest a hook when they depredate catch from longline hooks, which may become lodged in their mouth, throat, or stomach (Secchi *et al.* 2005, Fig. 3A). These events may lead to internal injuries, infections, starvation, or even eventual death (Best *et al.* 2001). Some hooked individuals may be unable to reach the surface to breathe, thus leading to a more immediate death by drowning (Hamer 2009a). Depredating odontocetes are also often conspicuous, especially during hauling when they are close to the vessel, which may lead to fishermen becoming frustrated and attempting to shoot individuals (Northridge and Hofman 1999). The impact of these mortalities at a population level is difficult to determine, because there are currently inadequate data available to estimate bycatch or mortality levels, or to determine historical and current levels (thus trends) in population size.

The distributions of most of the 72 extant odontocete species overlap geographically with longline fishing activities in some part of their range (Northridge 1984, Bjordal and Løkkeborg 1996, Culik 2004, Carwardine 2006). This is corroborated by the occurrence of operational interactions of 20 of the 72 extant odontocete species with longline fisheries (Table 1). The literature cited provides insights into this issue, although there is a much larger source from anecdotal, qualitative, unverified, and fishery dependent reports, indicating the occurrence of a much larger and chronic welfare and conservation problem.

A recent, intensive study of operational interactions between two odontocete species and a pelagic longline fishery in Hawaiian waters, based on independent longline observer programs and odontocete population surveys during the 1990s and 2000s, indicated that the decline of two odontocete populations may be attributable to bycatch mortalities on pelagic longline hooks (*e.g.*, pilot whale: Waring *et al.* 2006, Garrison 2007; false killer whale: Forney and Kobayashi 2007, Reeves *et al.* 2009). Nonetheless, establishing a robust quantitative link is difficult in the absence of reliable estimates of odontocete bycatch and populations, and of quantitative overlap, which is unlikely to occur with the data currently available (Hamer *et al.* 2008, 2009, Indian Ocean Tuna Commission 2010). This problem is further exacerbated by under reporting in fishery logbooks, which occurs because fishermen are typically fearful of the negative consequences of reporting accurately (Moore *et al.* 2010).

Recent advances in population genetics have made it possible to identify “management units,” which may assist in ensuring the biological importance of subpopulations is not underestimated (Pimper *et al.* 2010). Notwithstanding, most odontocete species have low reproductive rates and correspondingly low intrinsic capacities for increase, suggesting that even low levels of additional or unnatural mortality may cause decline (Leatherwood *et al.* 1983, Wade 2002, Culik 2004, Miller 2007). This is further complicated by the growing number of reports of genetic subdivision within what were previously thought to be single populations (*e.g.*, killer whale: Pilot *et al.* 2010; false killer whale: Chivers *et al.* 2007; common dolphin: Bilgmann *et al.* 2008; bottlenose dolphin: Krutzen *et al.* 2004). Nonetheless, this technology will allow more appropriate management strategies to be developed and implemented in the future, which take into account the genetic diversity *within* a species.

Although an individual odontocete is faced with the risk of injury or death when depredating from longlines, it may receive considerable foraging and energetic benefits by doing so. Some fish species caught on longlines may be unavailable to depredating odontocetes under natural conditions, because those fish are too large, too fast to catch, or occur in very deep waters (Gilman *et al.* 2006, Tixier *et al.* 2009). Fish




Figure 3. (A) False killer whale caught on a pelagic longline hook in Hawaiian waters (source: US National Marine Fisheries Service), (B) albacore depredated by odontocetes, with the torso completely removed from behind the gill plates, (C) odontocete tooth lacerations on torso of depredated albacore, (D) for comparison, damage caused by depredating shark, showing much cleaner removal of torso, (E) small bight marks on a sardine (*Sardinops sagax*) used for bait, probably caused by small depredating fish or squid, (F) large numbers of small fish in the vicinity of longline fishing gear that may be involved in bait depredation.

caught on longlines may offer an energetic advantage to depredating odontocetes, because they can be consumed without the need for deep dives or prolonged pursuits (Guinet *et al.* 2007). If an individual odontocete can develop a strategy to avoid becoming bycatch and the activities of the longline fishery they depredate from is frequent and predictable, then they may be at an energetic advantage compared with other individuals of the same species that forage naturally.

IMPACTS OF DEPREDAATION ON LONGLINE FISHERIES

Although concerns about the welfare and conservation of depredating odontocetes have become more common in recent times, concerns about the economic impact of depredation on affected longline fisheries have persisted since the 1960s (Dahlheim 1988, Yano and Dahlheim 1995, South Pacific Regional Environment Program 2002, Australian Fisheries Management Authority 2005, Indian Ocean Tuna Commission 2007). Depredation can reduce the overall size and condition of the landed catch, because target fish may be deterred from taking baited hooks, or caught fish may be damaged or removed completely (Yano and Dahlheim 1995, Northridge and Hofman 1999, Gilman *et al.* 2006, Hamer 2009*b*, Lauriano *et al.* 2009). When depredation occurs, affected fisheries are likely to experience sporadic, seasonal, or ongoing reductions in profit, which may lead to economic decline, especially if the problem persists.

When depredating odontocetes attack fish caught on longline hooks, they often remove the entire torso from behind the gill plates (Fig. 3B), or leave tooth lacerations on the torso (Fig. 3C). Odontocete teeth are pencil-like and tend to tear the skin and flesh of the caught fish, causing extensive damage. The nature of this damage is distinct from that caused by sharks, which tend to remove clean, bite-shaped portions of flesh from the torso of caught fish with their blade-like teeth, leaving the surrounding flesh relatively undamaged (Fig. 3D). Distinguishing between odontocete and shark depredation is important for (1) ensuring the correct attribution of damage to each depredating taxa and (2) selecting the correct mitigation method. Anecdotal accounts suggest that odontocetes may also be blamed for shark depredation. This is because depredating odontocetes surface frequently to breathe often in the vicinity of fishing vessels, thus are much more conspicuous than sharks. These instances may result in the overestimation of whale depredation and the underestimation of shark depredation.

In addition to impacting directly on the catch, depredating odontocetes may damage the fishing gear when they remove caught fish, specifically hooks, or larger portions of the longline gear, especially if they become caught themselves (Northridge and Hofman 1999, Gilman *et al.* 2006). Furthermore, small odontocetes (*i.e.*, dolphins) may partly (Fig. 3E) or completely remove baits directly from the hook (Secchi *et al.* 2005, McPherson *et al.* 2008). However, large quantities of small pelagic fish have been observed grazing on discarded baits and around baited hooks as they are hauled aboard the vessel at the end of a set (Hamer, personal observation; Fig. 3F). This suggests small pelagic fish or squids, rather than odontocetes, may be responsible for depredating baits on some occasions.

The damage to caught fish and deterrence of target fish by depredating toothed whales is likely to result in financial losses to affected fisheries. Two studies conducted in the Bering Sea, northeast Pacific, between 1977 and 1989 estimated the daily economic cost of killer whale depredation to each vessel across a Greenland demersal

fishery (for halibut, *Reinhardtius hippoglossoides*, and arrowtooth flounder, *Atheresthes stomias*) was between US\$1,034 and US\$8,449 (Dahlheim 1988, Yano and Dahlheim 1995). These figures were based on one set per day and 78.9% inflation between 1989 and 2010 for the earlier study and 44.2% between 1995 and 2010 for the later study. Two other studies conducted around the Crozet (46°25'S, 50°59'E) and Kerguelen Islands (49°19'S, 69°28'E), Southern Ocean, between 2003 and 2008 estimated the daily cost of killer whale and sperm whale depredation to the fishery across a French demersal fishery (for Patagonian toothfish, *Dissostichus eleginoides*) was between US\$6,052 and US\$8,495 (Roche *et al.* 2007, Tixier *et al.* 2009). These figures were adjusted from a multiyear to a one set per day estimate, then 1.5:1€ to US\$ exchange conversion and 13.6% inflation between 2005 and 2010 for the earlier study and a 1.3:1€ to US\$ exchange conversion and 2.7% inflation between 2008 and 2010 for the later study. It is important to note that these figures are at best informative and represent a snapshot in time for two demersal longline fisheries in the northeast Pacific and the Southern Ocean. They are unlikely to reflect the losses sustained by other demersal or pelagic longline fisheries in other locations at other times. In addition, neither study reported the overall catch figures for vessels or for the fleet, thus it was not possible to determine the percentage of catch that was lost, nor the impact on profits. Furthermore, the economic cost of depredation is likely to be an underestimate, because it is not possible to quantify the number of caught fish that are completely removed from the hook, the number of target fish that are deterred from taking a baited hook altogether, the amount of fishing gear that is damaged, nor the various avoidance activities undertaken by skippers (Yano and Dahlheim 1995, Hamer 2009*b*). Despite the lack of data, the economic costs reported in the two studies detailed suggest that the fishery wide economic impact is likely to be significant.

When depredating whales completely remove caught fish from longline hooks, they may impede effective fishery management practices, such as setting accurate total allowable commercial catch (TACC) limits and calculating the catch per unit of effort (CPUE) for target fish stocks. Overfishing may occur, because the caught fish that are removed by depredating odontocetes are not included when calculating the level of exploitation of the targeted fish (Gilman *et al.* 2006). This situation may occur when the TACC has already been set, because the depredated fish are not included in the catch declared by the fishery. Under fishing may occur, because the removal of caught fish by depredating whales will reduce the CPUE. This situation may lead to the false impression that there are less fish available than is actually the case, which may encourage the relevant fishery management agencies to become cautious and thus reduce the TACC (Gilman *et al.* 2006, Roche *et al.* 2007). Therefore, the impact that catch depredation by odontocetes has on the management of the affected fishery should be taken into account when determining methods for long term sustainable fishery management.

BYCATCH AND DEPREDAATION MITIGATION STRATEGIES

Until recently, few studies have attempted to identify solutions for mitigating operational interactions between odontocetes and longline fishing operations. Most simply flagged promising ideas (*e.g.*, Northridge and Hofman 1999, Visser 2000), while some compiled more detailed accounts of mitigation measures implemented directly by individual longline fishermen and by fisheries in an *ad hoc* and untested

manner (*e.g.*, Dahlheim 1988, Secchi *et al.* 2005, Table 2). Independent experimental trials that quantify the effectiveness of potential depredation mitigation strategies in the longline industry remain in their infancy (*e.g.*, Moreno *et al.* 2008, Mooney *et al.* 2009). In contrast, the available literature indicates that the development and implementation of cetacean depredation and bycatch mitigation strategies in other fisheries is more advanced (*e.g.*, purse seine: Gosliner 1999, Hamer *et al.* 2008; gill net: Trippel *et al.* 1999, Barlow and Cameron 2003).

Fishermen, fishery managers, and researchers have proposed a number of techniques for mitigating odontocete depredation and bycatch in longline fishing gear, which can be broadly categorized as (1) behavioral, (2) spatial, (3) acoustic, or (4) physical. Behavioral techniques have been successfully used in active fishing methods, such as trawl (Tilzey *et al.* 2006) and purse seine (Hamer *et al.* 2008), because the fishing gear can be manipulated and monitored during a fishing event to mitigate the likelihood of bycatch. Unfortunately, there is limited scope for the behavior of fishermen to influence the way longline gear behaves when it is deployed, because it hangs passively in the water column, is not attached to the vessel, and is generally remote and out of sight. Nonetheless, fishermen are able to make decisions about where to fish and for how long to deploy the gear in order to avoid odontocetes, although such practices are yet to be quantified and are unlikely to be implemented voluntarily, especially in the long-term, unless the economic benefits are immediately apparent. The “move on” tactic has been used by some longline fishermen in a bid to outrun depredating whales, although the success of this strategy seems to be ambiguous at best and is likely to be costly, thus affecting profit margins.¹ A study of this method for avoiding pinniped depredation and bycatch in a trawl fishery found it was only occasionally successful, because depredating individuals were also able to travel long distances to remain with the vessel (Tilzey *et al.* 2006).

Spatial closures, typically known as marine protected areas (MPAs), are designed to spatially separate marine mammal populations and fishing effort so as to reduce the likelihood of bycatch mortalities. However, MPAs that are effective in protecting odontocete populations are difficult to implement, because (1) knowing where to put them is often difficult to determine in the absence of reliable data on odontocete migration and movement patterns (Dulau-Drouot *et al.* 2008), (2) they are often smaller than necessary, due to stakeholder pressure to minimize their impact on fisheries (Klein *et al.* 2008), (3) monitoring compliance by fishermen is difficult due to the lack of capacity and resources (Le Quesne 2009) and (4) quantitative performance assessment is hampered by the statistical uncertainties associated with limited and potentially unrepresentative data (Claudet *et al.* 2006). In contrast, the implementation of MPAs to protect pinnipeds from fishing activities has proven easier, because they are central place foragers whose at-sea movements and population trends can be determined with comparative ease (Baylis *et al.* 2008, Shaughnessy *et al.* 2011, Hamer *et al.* 2011).

Acoustic Technologies

Acoustic technologies for mitigating odontocete depredation and bycatch in longline fisheries have received the most attention in the literature. In general, high intensity sounds are used to deter depredating odontocetes from approaching fishing

¹Personal communications from Mark Coker and Niki McCulloch, De Brett Seafood, Queensland, Australia, November 2009 and June 2011, respectively.

Table 2. Summary of methods previously considered or trialed by fishermen and researchers to mitigated catch depredation by whales from longlines.

Category and type	Method		Result Success/failure	Problems		Source
	Description			Realized or perceived		
Physical Net sleeve	Branch line mounted. Prevents access. Passively drops over hooks and caught fish during hauling.		Success ^a	Intelligent animals have learned to damage tail of fish	1	1
			Success ^{a,b}			
Metal wire	Line mounted. Flaps about to deter cetacean. Descends troll line when fish is caught.		Pending outcome ^a	Dependent on whales being deterred by the presence of streamers.	3, 7, 8	3, 7, 8
Streamers/tangles	Snood mounted. Flaps about to deter cetacean. Descends snood when fish caught.			Requires complex device, so may have maintenance problems.		
Chemical Lithium chloride/ether	Elicits vomit response. Mounted near hook. Activated when fish caught.		Not trialed	Unknown health issues for depredating whales and humans. Potential ethical issues.	3, 4, 8	3, 4, 8
Stress/decay marker	Elicits escape/exit response. Mounted near hook. Activated when fish caught.		Not trialed	May dissipate too quickly, or have adverse effects over wide area.	4, 8	4, 8
Electrical Stinger	Snood mounted. Deployed when fish caught and activated when cetacean approaches.		Pending outcome ^a	Potential ethical issues for cetaceans and safety issues for crew.	3, 4	3, 4

Continued

Table 2. (Continued)

Category and type	Method	Description	Result Success/failure	Problems Realized or perceived	Source
Visual Bubble screen	Interferes with visual sense.		Not trialed	May be difficult to maintain.	
Acoustic Detection	Use of listening devices to pick up echolocation signals from cetaceans in the area.		Limited success ^a	Logistically difficult to achieve over wide area. Results are often ambiguous and inconclusive.	4, 7
Predator playback	Use of predator noises to elicit escape response such as killer whale calls to deter pilot whales.		Not trialed	Works over an insufficient distance. Individuals may become habituated, making them vulnerable.	4
Masking/disruption	Producing predominant "white noise" to mask noises produced by vessel activities.		Initial success	Works over insufficient distance. Trialed on a captive animal only.	4, 6
Harassment	Annoying and potentially damaging sound forces cetaceans to leave the area.		Unsuccessful ^a	Demonstrated learning by individual reduced device performance. May cause hearing damage and stranding.	4
Accessory skills	Acoustic novelty draws cetaceans away from fishing gear.		Not trialed	May have adverse effects on other animals. Would only work on demersal longlines where line comes up to boat. Logistically difficult to achieve for pelagic longlines.	4

Quiet operations	Modify vessels to make less noise.	Initial success	Individuals may learn to detect signatures in background noise.	3, 5, 8
Explosives/seal bombs	Loud noise causes flight response.	Unsuccessful	May cause hearing damage and stranding. May have adverse effects on other animals.	4
Behavioral Operant conditioning	Behavioral modification using signal cues.	Not trialed	Requires high proportion of animals in the population to learn.	4
Blank sets	Gear set without baits to confuse whales.	Unsuccessful	Depredating individuals soon learned to search for baited sets.	
Management Spatial closures	Away from areas frequented by depredating cetaceans.	Not trialed	Moves effort to a different location—may cause other problems. Often puts effort outside prime fishing ground.	7, 8
Temporal closures	Away from areas frequented by depredating cetaceans at certain times of the year.	Not trialed	Moves effort to a different time of year—may cause other problems.	7, 8
Move fishery	Away from traditional fishing grounds to areas not frequented by depredating cetaceans.	Limited success	Often puts effort outside prime fishing period. Large volume of fuel to move >60 nmi.	4, 9
			Often puts vessels outside prime fishing ground.	

Continued

Table 2. (Continued)

Category and type	Method		Result	Problems		Source
	Description	Success/failure		Realized or perceived		
Change target species	To a species thought to be unattractive to depredating cetaceans.	Mixed results	Alternative species often more difficult to catch or less valuable.	4, 8		
Change time of fishing	Fish at night instead of during the day.	Unsuccessful	Depredating whales learn to take advantage of new food source.	4		
Change depth of set	Out of depth range of depredating cetaceans.	Limited success	May only be effective for species that only feed during the day.	8, 9		
Change gear type	Use pots to catch the fish instead of longlines.	Limited success	May also put gear beyond depth of target fish species.	4, 7		
Culling	Shooting or harvesting of cetaceans.	Not trialed	Possible only in demersal fisheries. Often results in reduced catch. Illegal and unethical.	4		

Information source:

- 1 Moreno *et al.* 2008
- 2 Zollert and Read 2006^b
- 3 Hamer 2009^b
- 4 Dahlheim 1988
- 5 AFMA 2005
- 6 Mooney *et al.* 2009
- 7 McPherson *et al.* 2008
- 8 Gilman *et al.* 2006
- 9 Tixier *et al.* 2009

^aOutcome based on experimental trials.

^bStudy of a troll fishery—included here due to the relevance of the depredation mitigation strategy to longline fishing.

gear, while comparatively low intensity sounds are used to alert odontocetes (and other cetaceans) to the presence of fishing gear to prevent them from becoming incidental bycatch.

The majority of the literature in this field reports on ways to mitigate catch depredation and typically focus on four strategies, which are harassment, deterrence, echolocation disruption, and avoidance. Their development has generally been encouraged by fishery stakeholders who hope to reduce the economic impact of depredation on their fishing enterprise. Most effort has focused on the development of acoustic harassment devices (AHDs), which are designed to encourage or force depredating odontocetes to leave the vicinity of the fishing gear (Nowacek *et al.* 2007). In the absence of information on odontocete hearing range and capacity, AHD development seems to have been based on human characteristics. As such, they typically transmit sounds greater than 180 dB (at 1 m from the source), which are beyond the 145 dB level at which human hearing structures are at risk of permanent damage (Price 1981, Nowacek *et al.* 2007).

Acoustic deterrence devices (ADDs) emit moderately high sounds that are typically lower than 180 dB (at 1 m from the source). Based on similar principals to AHDs, ADDs are designed to annoy depredating odontocetes and encourage them to leave or remain clear of an area, typically where floating structures such as fish pens are located (Dawson *et al.* 1998, Nowacek *et al.* 2007). Unlike AHDs and ADDs, echolocation disruption devices (EDDs) are claimed to prevent depredating odontocetes from accurately determining the location of a caught fish, thus reducing the likelihood of a successful depredation attempt (Mooney *et al.* 2009). Nonetheless, the distinction between ADDs and EDDs remains unclear, because the functional mechanisms that elicit specific behavioral outcomes in depredating odontocetes in response to the sound emitted remains unclear (Jefferson and Curry 1996).

Efforts have also been made to avoid depredation altogether. Passive listening arrays (PLAs) are designed to assist affected fishing vessels in acoustically detecting depredating odontocetes, thus allowing the vessel to leave or move on from a fishing area when individuals are detected in the vicinity (McPherson *et al.* 2008). Unlike AHDs, ADDs, and EDDs, PLAs are not a deterrent mechanism, instead providing fishermen with the ability to reduce the level of physical overlap with depredating odontocetes. Despite the apparent benefits, this technology remains in its infancy. This is partly due to the difficulties associated with confirming the presence or absence of highly mobile odontocetes, with depredation occurring at times when odontocetes are not observed or detected and depredation not occurring at times when odontocetes are observed or detected.² In addition, PLAs have been slow to develop, because the associated structures and equipment are typically complex and expensive (Nielsen and Møhl 2006), they can be damaged by marine predators such as sharks (Johnson *et al.* 1982) and sound interference from the fishing vessel and from the broader marine environment can mask the vocalizations of depredating whales (Thode *et al.* 2007).

An alternative strategy for avoiding depredating odontocetes involves masking or minimizing the sound signature of fishing vessels that are thought to attract depredating odontocetes, so that the individuals involved can no longer detect the vessels presence (McPherson *et al.* 2008). However, identifying a parsimonious suite

²Personal communication from Mark Coker, De Brett Seafood, Queensland, Australia, November 2009.

of sound signature and suppression factors that may assist in mitigating depredation has proven to be logistically and technically challenging.³

Acoustic technology has also been used to mitigate incidental bycatch of odontocetes, with the main strategy being to warn individuals of the presence of fishing gear in their vicinity (Kraus *et al.* 1997, Barlow and Cameron 2003). Such devices are referred to as “pingers.” Their development and implementation has predominantly occurred in association with gill net and drift net fisheries, where incidental bycatch of odontocetes seems to have been greatest (Read *et al.* 2006). A number of studies have shown that pingers significantly reduce incidental bycatch of harbor porpoises (*Phocoena phocoena*) in demersal gill nets (Lien *et al.* 1995, Kraus *et al.* 1997, Trippel *et al.* 1999, Gearin *et al.* 2000) and common dolphins in drift gill nets (Barlow and Cameron 2003). However, other studies have shown that assessing the effectiveness of pingers is difficult, due to the lack of statistical power caused by typically low rates of bycatch, habituation of individuals to the noises made by pingers (which can also lead to the ‘dinner bell effect’, where individuals are actually attracted by the noise emitted, rather than being warned away or repelled) and a lack of understanding of the processes that lead to bycatch (Dawson *et al.* 1998). Pingers may also be species specific in their application, thus making it difficult to address bycatch effectively, especially in situations where more than one species is involved (Kastelein *et al.* 2006). Despite the potential usefulness of pingers in some fisheries, they are unlikely to be useful in longline fisheries, where most bycatch occurs when depredating odontocetes are actively attempting to remove caught fish from hooks.

Despite the considerable efforts to develop acoustic strategies for mitigating depredation and bycatch, to date their successful application has proven difficult. The use of AHDs raises ethical concerns about the effect of high-level noise on odontocetes and the wider marine environment (Johnston and Woodley 1998, Morton and Symonds 2002). One study of the use of ADDs in a gill net fishery suggested that the subsequent reduction in dolphin bycatch may have occurred because the devices deterred the target fish, thus encouraging the dolphins involved to forage elsewhere (Kraus *et al.* 1997). Over time, odontocetes may become habituated to noises emitted by ADDs, EDDs and pingers, thus rendering them ineffective (Jefferson and Curry 1996). These devices may eventually become attractants or a “dinner bell” to foraging odontocetes, if animals learn to associate the sounds emitted with the presence of palatable fish (Jefferson and Curry 1996, Mooney *et al.* 2009).

Assessing the efficacy of acoustic devices has been hampered by the lack of experimental replication, mainly due to the variety of odontocete species involved, the number of devices currently available in the market place, variations in the configuration of gear used in each fishery and the unique environmental conditions in each coastal or oceanic region (Jefferson and Curry 1996, Dawson *et al.* 1998, Nowacek *et al.* 2007, Kastelein *et al.* 2006). Current technology also constrains the testing and application of acoustic devices, because integral components such as the transponder and batteries are currently large and expensive (McPherson *et al.* 2008, Mooney *et al.* 2009). These problems are further exacerbated by the current lack of understanding of the mechanisms that underpin how noise harasses, deters or warns odontocetes that are close to fishing gear. While there is a need to continue developing and assessing acoustic depredation and bycatch mitigation strategies in longline fisheries, success, and ultimately implementation, will only be possible if a case-by-case approach is adopted, experiments are controlled and replicated, and the necessary components

³Personal communications from Mark Coker, De Brett Seafood, Queensland, Australia, November 2009; Will Mure, Mures Fishing, Tasmania, Australia, January 2010.

are sufficiently small and cheap for devices to be deployed in large numbers on the gear.

Physical Technologies

Physical technologies for mitigating depredation and bycatch of odontocetes have received comparatively little attention. However, recent innovations in developing physical depredation mitigation devices (PDMDs) have proven promising in a Chilean demersal longline fishery for Patagonian toothfish, with the experimental testing of the 'net sleeve' demonstrating that sperm whale depredation could be reduced by 82.8% (Moreno *et al.* 2008, Hamer 2010). Typically, odontocetes are unable to depredate caught fish when demersal gear is deployed on the benthos, with access only possible during the latter stages of the haul. The rigid net sleeve was designed to remain clear of the baited hooks on the benthos during fishing, then descended the branchline under the influence of gravity, thus preventing access to the caught fish by depredating sperm whales (Fig. 4).

Developing solutions for pelagic longline gear is likely to be more challenging than for demersal longline gear. Pelagic longlines are set at much shallower depths (*i.e.*, between 30 m and 300 m from the surface), thus are accessible by most depredating odontocete species throughout the fishing period (Hamer 2010). As such, an effective device would need to be comparatively complex, remaining clear of the baited hook to allow it to function unimpeded. The device must also include a trigger mechanism that is activated by line tension when a target fish becomes caught and attempts to escape. Efforts to solve this problem in other hook-based pelagic fisheries may provide valuable insights. A recent study reported on attempts to mitigate catch depredation




Figure 4. Schematic 'time-lapse' view of net sleeve operation, showing the protection of the fish and the physical deterrence of a sperm whale during the haul (With permission: Carlos A. Moreno).

by bottlenose dolphins (*Tursiops truncatus*) in the Florida king mackerel (*Scomberomorus cavalla*) troll fishery and found that dolphins were deterred from depredating by a “metal wire” that moved around in the water next to the caught fish (Zollett and Read 2006). In order to allow the baited hook to fish unimpeded, the metal wire was held clear in a tension sensitive mechanism. When a fish was caught and the tension increased, the metal wire was released and then descended the line towards the caught fish. It was assumed that dolphins were deterred from depredating the caught fish due to fear of physical injury or entanglement.

Although the development of PDMDs (to mitigate both bycatch of and depredation by odontocetes) is in its infancy, they offer a comparatively realistic, applicable and generic approach when compared with the current generation of available acoustic devices. This is because it is possible to manufacture comparatively small and cheap devices that can be placed on each snood immediately adjacent the hook, where depredation and bycatch events take place. Testing their efficacy is also comparatively simple, with the use of rigorous and controlled experimental trials that measure target fish catch rates, and odontocete depredation and bycatch rates. In contrast, the efficacy of acoustic devices is more difficult to determine, because it is not possible to ascertain whether the presence of a device is directly responsible for the results obtained.

Despite the conceptual and experimental advantages associated with developing PDMDs, accommodating the necessary physical complexity of the devices used in pelagic longline fisheries may prove challenging. Unlike acoustic devices that generally have no external moving parts, PDMDs may contain many. A conventional pelagic longline operation involves the deployment of between 1,800 and 3,600 hooks, with a hook deployed every 6–8 s during setting, indicating that the addition of the few extra seconds needed between each hook to attach or remove PDMDs may considerably increase the duration of daily fishing activities (Hamer 2009*b*). In addition, PDMDs need to be small and complex, which may reduce their durability in the harsh marine environment.

Pelagic longliners targeting tuna and billfish in the Tropical South Pacific Ocean (TSPO) have unofficially reported depredation by pilot whales and false killer whales for over a decade. In more recent times, a number of anecdotal reports by fishermen have indicated that depredating individuals may be avoiding sections of the longline where the gear has become tangled, mainly as a result of vigorous and prolonged swimming activity by caught billfish and sharks.⁴ As such, an approach reminiscent of the Florida king mackerel example may be worthwhile, where structures that simulate or mimic tangles are deployed near a caught fish to both physically and psychologically deter depredating odontocetes. The Australian Government has initiated a study of this nature, which is currently in the developmental phase, although extensive sea trials on commercial longline vessels will soon commence in the TSPO and in the Indian Ocean (Hamer 2011, Fig. 5).

The success of PDMDs is dependent on the level of industry implementation, which is more likely to be associated with voluntary uptake, rather than mandatory or enforcement means. Mandating or enforcing the use of PDMDs that have no obvious benefit to the fishery would likely result in the need for costly monitoring activities and in some level of noncompliance, both of which are unfavorable outcomes.

⁴Personal communications from Mark Coker, De Brett Seafood, Queensland, Australia, November 2009; Ueta Faasili, Samoa Ministry of Agriculture and Fisheries, Samoa, February 2010; Tom Mayo, Solander Fisheries, Suva, Fiji, August 2011.




Figure 5. Schematic diagram of the *chain device* and of the *cage device* (A, C: not triggered; B, D: triggered) currently under development by the Australian Government and soon to be trialed in the Pacific and Indian Oceans. Before the devices are triggered by the tension of a caught fish, they remain clear of the baited hook and close to the mainline or swivel. Upon being triggered, the devices release the streamers or cage and then descend the snood toward the caught fish, eventually enveloping it.

Specifically, fishermen are unlikely to purchase and implement this technology if the cost of doing so is more than the increases in revenue associated with reduced catch damage, suggesting the aim should be to ensure a cost-benefit analysis will work in the favor of the fisherman. This is especially important when considering the large number of illegal, unregulated, and unreported (IUU) pelagic longline fisheries around the world that avoid conventional management regulations (Food and Agriculture Organization 2001, Baker *et al.* 2006, Lukoschek *et al.* 2009). As

such, PDMD development should not only focus on efficacy and durability, but also on cost minimisation.

SUMMARY AND FUTURE DIRECTIONS

The literature summarized here indicates that odontocete bycatch and depredation in longline fisheries is widespread, involving many fisheries and many odontocete species and populations. Mitigating this problem is becoming a higher priority for all stakeholders, especially as longline fishery profit margins dwindle and as competition and conflict between odontocetes and longline fisheries increase. Nonetheless, the problem of depredation may be overstated by some fishermen who attribute poor catch performance to odontocete depredation, when the real cause may be poor operational decisions and thus poor catch performance, or incorrect assignment of depredation by other taxa (*i.e.*, sharks, fish or squid). In contrast, the problem may be understated for odontocetes, because small and as yet unidentified populations that include depredating individuals may be at risk of decline with the loss of only a few individuals (Leatherwood *et al.* 1983, Beissinger and McCullough 2002, Miller 2007). An earlier review of this issue indicated that the successful mitigation of bycatch and depredation can only occur with changes in longline fishing practices, although it lamented that little had been done thus far to identify and test potential solutions (Gilman *et al.* 2006). Given the volume of information summarized here, the geographic extent of operational interactions and the efforts made to develop mitigation strategies to date, stakeholders are strongly encouraged to prioritize this problem without delay, with a view to conserving affected odontocete populations and sustaining longline fisheries.

Both acoustic and physical mitigation technologies have shown some promise, although both appear have inherent problems that may hinder their development and implementation. Acoustic mitigation tools are simple in their application, although cumbersome and inadequate in their function and complex in their assessment. For example, the size of the batteries and transponders currently available are probably too big for most applications, while the types and levels of noises that deter specific odontocete species remain unclear and testing the efficacy of those noises in a highly dynamic marine environment is difficult. By comparison, PDMDs are simple in their function and assessment, although their practical application and implementation may be challenging if maintenance and per-unit cost are high. Perhaps the most important aspect of ensuring the success of any bycatch and depredation mitigation strategy, whether it be acoustic, physical, or any other form, is the need to keep purchase and implementation costs below the economic gains associated with increased catch revenue. Although some fishery management agencies may opt for mandatory implementation of these technologies, the reality is that noncompliance is likely to be widespread if there are no economic benefits for the affected fishery. Longline fishermen involved in IUU activities are likely to monitor developments in regulated or managed fisheries and will only adopt mitigation technologies if there is a perceived economic benefit. Therefore, stakeholders involved in the development of PDMDs and other depredation and bycatch mitigation technologies should aim to minimize costs in order to increase the likelihood of voluntary implementation by the affected fishery regardless of its management status.

Individual longline operations and fisheries around the globe are likely to face a diversity of situations, such as differences in (1) the odontocete species they interact with and bycatch rates, (2) target fish depredation rates, catch rates and value, and (3)

overall operational costs (including repayments, fuel, wages, bait, *etc.*). Given that a single panacea to this diverse problem is unlikely to emerge, fishermen and fishery managers are encouraged to maximize the chance of mitigating odontocete bycatch and depredation by using a suite or “toolbox” of mitigation strategies, such as (1) acoustic, (2) physical, (3) fishermen behavior, and (4) MPAs (*e.g.*, Dahlheim 1988, Gilman *et al.* 2002, Gilman *et al.* 2006, Campbell and Cornwall 2008).

The development and implementation of acoustic and physical mitigation methods has attracted the interest of a broad stakeholder base. At the policy end of the spectrum, regional partnerships and agreements are deemed necessary for facilitating necessary research and for securing necessary funds. For example, the joint tuna regional fishery management organizations (T-RFMOs) have implemented the “Kobe bycatch process” to deal with issues of whale bycatch and depredation. However, a number of delegates at the *Kobe II Bycatch Workshop* (Brisbane, Australia) in 2010 criticized the process, citing (1) a lack of consensus about fundamental terminology (*e.g.*, what constitutes bycatch?), (2) the overcomplicated and slow process, and (3) the misguided focus on documentation rather than problem solving (Kobe II 2010). At the operational end of the spectrum, researchers appear to have made extensive inroads toward finding solutions, by focusing a growing body of knowledge on the subject of mitigating odontocete bycatch and depredation (*e.g.*, Gilman *et al.* 2006, Zollett and Read 2006, Moreno *et al.* 2008, McPherson *et al.* 2008, Hamer 2011). Nonetheless, researchers have acknowledged that the conceptualization and development of some solutions may lie with the fishermen whose knowledge, experience and enthusiasm should not be underestimated or undervalued (Gilman *et al.* 2006). Regardless of where one might sit on the policy and operations spectrum, it is clear that greater resources and commitment need to be focused on assisting fishermen and researchers to address this issue at the vessel, preferably at the hook. If this does not occur soon, it should be expected that operational interactions (depredation and bycatch) between odontocetes and longline fisheries will continue or increase, thus casting uncertainty on the future of affected longline fisheries and on the fate of affected odontocete populations.

LITERATURE CITED

- Ashford, J. R., P. S. Rubilar and A. R. Martin. 1996. Interactions between cetaceans and longline fishery operations around South Georgia. *Marine Mammal Science* 12:452–457.
- Australian Fisheries Management Authority (AFMA). 2005. Mammal depredation on demersal longlines: A review prepared by AFMA for the gillnet, hook and trap fishery. Report, Australian Fisheries Management Authority (AFMA). April 2005. 24 pp.
- Baker, C. S., V. Lukoschek, S. Lavery, M. L. Dalebout, M. Yong-Un, T. Endo and N. Funahashi. 2006. Incomplete reporting of whale, dolphin and porpoise ‘bycatch’ revealed by molecular monitoring of Korean markets. *Animal Conservation* 9:474–482.
- Bakun, A., E. A. Babcock and C. Santora. 2009. Regulating a complex adaptive system via its wasp-waist: Grappling with ecosystem-based management of the New England herring fishery. *Journal of Marine Science* 66:1768–1775.
- Barlow, J., and G. A. Cameron. 2003. Field experiments show that acoustic pingers reduce marine mammal bycatch in the California drift gill net fishery. *Marine Mammal Science* 9:265–283.
- Baylis, A. M. M., B. Page and S. D. Goldsworthy. 2008. Colony-specific foraging areas of lactating New Zealand fur seals. *Marine Ecology Progress Series* 361:279–290.

- Beisinger, S. R., and D. R. McCullough. 2002. Population viability analysis. The University of Chicago Press, Chicago, IL.
- Bell, C., P. D. Shaughnessy, M. Morrice and B. Stanley. 2006. Marine mammals and Japanese longlining fishing vessels in Australian waters: Operational interactions and sightings. *Pacific Conservation Biology* 12:31–39.
- Best, P. B., V. M. Peddemors, V. G. Cockcroft and N. Rice. 2001. Mortalities of right whales and related anthropogenic factors in South African waters, 1963–1998. *Journal of Cetacean Research and Management* 2:171–176.
- Beverton, R. J. H. 1985. Analysis of marine mammal—fisheries interactions. Pages 3–33 in J. R. Beddington, R. J. H. Beverton and D. V. Lavigne, eds. *Marine mammals and fisheries*. George Allen and Unwin, London, U.K.
- Bilgmann, K., L. M. Moller, R. G. Harcourt, R. Gales and L. B. Beheregary. 2008. Common dolphins subject to fisheries impacts in southern Australian are genetically differentiated: Implications for conservation. *Animal Conservation* 11:518–528.
- Bjordal, Å., and S. Løkkeborg. 1996. Longlining. Fishing News Books, Oxford, U.K.
- Campbell, L. M., and M. L. Cornwall. 2008. Human dimensions of bycatch reduction technology: Current assumptions and directions for future research. *Endangered Species Research* 5:325–234.
- Carwardine, M. 2006. Whales, dolphins and porpoises. 2nd edition. Dorling Kindersley, London, U.K.
- Chivers, S. J., R. W. Baird, D. J. McSweeney, D. L. Webster, N. M. Hedrick and J. C. Salinas. 2007. Genetic variation and evidence for population structure in eastern North Pacific false killer whales (*Pseudorca crassidens*). *Canadian Journal of Zoology* 85:783–794.
- Claudet, J., D. Pelletier, J. Y. Jouvenel, F. Bachet and R. Galzin. 2006. Assessing the effects of marine protected area (MPA) on a reef fish assemblage in a northwest Mediterranean marine reserve: Identifying community-based indicators. *Biological Conservation* 130:349–369.
- Culik, B. M. 2004. Review of small cetaceans: Distribution, behaviour, migration and threats. Report, United Nations Environment Program (UNEP), Convention on Migratory Species (CMS), Bonn, Germany. 84 pp.
- Dahlheim, M. E. 1988. Killer whale (*Orcinus orca*) depredation on longline catches of sablefish (*Anoplopoma fimbria*) in Alaskan waters. Report, Northwest and Alaska Fisheries Center (NWAFC), National Marine Fisheries Service (NMFS), 88–14. 38 pp. Available at <http://www.afsc.noaa.gov/Publications/ProcRpt/PR%2088-14.pdf> (accessed 30 July 2011).
- Dalla Rosa, L., and E. R. Secchi. 2007. Killer whale (*Orcinus orca*) interactions with the tuna longline fishery off southern and south-eastern Brazil: A comparison with shark interactions. *Journal of the Marine Biological Association of the United Kingdom* 87:135–140.
- Dawson, S. M., A. Read and E. Slooten. 1998. Pingers, porpoises and power: Uncertainties with using pingers to reduce bycatch of small cetaceans. *Biological Conservation* 84:141–146.
- Di Natale, A., and A. Mangano. 1983. Biological and distribution new data on the sperm whale, *Physeter macrocephalus* L., in the central Mediterranean Sea. Report, International Council for the Scientific Exploration of the Mediterranean Sea 28:183–184.
- Donoghue, M., R. Reeves and G. Stone. 2003. Report of the workshop: Cetacean interactions with commercial longline fisheries in the South Pacific region: Approaches to mitigation. Apia, Samoa, 11–15 November 2002. New England Aquarium Press, Boston, MA. 44 pp. Available at <http://www.nmfs.noaa.gov/pr/pdfs/interactions/samoa2002.pdf> (accessed 30 July 2011).
- Duarte, C. M., M. Holmer, Y. Olsen, *et al.* 2009. Will the oceans help feed humanity? *Bioscience* 59:968–976.
- Duguy R., and E. Husenot. 1982. Occasional captures of delphinids in the Northeast Atlantic. Report of the International Whaling Commission 32:461–462.

- Dulau-Drouot, V., V. Boucaud and B. Rota. 2008. Cetacean diversity off La Reunion Island (France). *Journal of the Marine Biological Association of the United Kingdom* 88:1263–1272.
- Earle, M. 1996. Ecological interactions between cetaceans and fisheries. Pages 167–204 in M. P. Simmonds and J. D. Hutchison, eds. *The conservation of whales and dolphins: Science and practice*. John Wiley and Sons, Chichester, U.K.
- Ebert, S., G. P. Metschies, D. Schmid and A. Wagner. 2009. *International fuel prices*. 6th edition. Report, Transport Policy Advisory Services, Federal Ministry for Economic Cooperation and Development, Eschborn, Germany.
- Food and Agriculture Organization. 2001. International plan of action to prevent, deter and eliminate illegal, unreported and unregulated fishing. Report, United Nations (UN) Food and Agriculture Organization (FAO), Rome, Italy. 24 pp. Available at <http://www.fao.org/docrep/003/y1224e/y1224e00.htm> (accessed 30 July 2011).
- Food and Agriculture Organization. 2009. *The state of world fisheries and aquaculture 2008*. Report, United Nations (UN) Food and Agriculture Organization (FAO), Rome, Italy. 196 pp. Available at <ftp://ftp.fao.org/docrep/fao/011/i0250e/i0250e.pdf> (accessed 30 July 2011).
- Forney, K. A., and D. Kobayashi. 2007. Updated estimates of mortality and injury of cetaceans in the Hawaii-based longline fishery, 1994–2005. NOAA Technical Memorandum NMFS-SWFSC-412. 30 pp.
- Garrison, L. 2007. Interactions between marine mammals and pelagic longline fishing gear in the U.S. Atlantic Ocean between 1992 and 2004. *Fishery Bulletin* 105:408–417.
- Gearin, P. J., M. E. Goshko, J. L. Laake, L. Cooke, R. L. Delong and K. M. Hughes. 2000. Experimental testing of acoustic alarms (pingers) to reduce bycatch of harbour porpoise, *Phocoena phocoena*, in the state of Washington. *Journal of Cetacean Research and Management* 2:1–10.
- Gilman, E., P. Gandini, C. Carboneras, G. Balogh, D. Leadbitter, and T. Neves. 2002. Incentive instruments for sustainable bycatch of sensitive species in longline fisheries. Proceedings of the Incentives Session, Second International Fishers' Forum, 19–22 November 2002. United States Western Pacific Regional Fishery Management Council, Honolulu, HI. 124 pp.
- Gilman, E., N. Brothers, G. McPherson and P. Dalzell. 2006. A review of cetacean interactions with longline gear. *Journal of Cetacean Research and Management* 8:215–223.
- Gosliner, M. L. 1999. The tuna-dolphin controversy. Pages 120–155 in J. R. Twiss and R.R. Reeves, eds. *Conservation and management of marine mammals*. Melbourne University Press, Melbourne, Australia.
- Guinet, C., P. Domenici, R. de Stephanis, L. Barrett-Lennard, J. K. B. Ford and P. Verborgh. 2007. Killer whale predation on bluefin tuna: Exploring the hypothesis of the endurance-exhaustion technique. *Marine Ecology Progress Series* 347:111–119.
- Hamer, D. J. 2009a. Update #1—October 2009: Whale-catch depredation mitigation project commences. Non-lethal options for mitigating catch depredation by toothed whales from pelagic longlines. pp. 1–4. Available at http://www.marinemammals.gov.au/__data/assets/pdf_file/0016/2518/1.-Update-Project-Background-and-Objectives.pdf (accessed 30 July 2011).
- Hamer, D. J. 2009b. Update #2—December 2009: First steps toward mitigating catch depredation by whales. Non-lethal options for mitigating catch depredation by toothed whales from pelagic longlines. pp. 5–18. Available at http://www.marinemammals.gov.au/__data/assets/pdf_file/0014/2930/2.-Update-Description-of-Problem-and-Possible-Solutions.pdf (accessed 30 July 2011).
- Hamer, D. J. 2010. Update #6—July 2010: PDMD progress. Non-lethal options for mitigating catch depredation by toothed whales from pelagic longlines. pp. 30–33. Available at http://www.marinemammals.gov.au/__data/assets/pdf_file/0018/29304/6.-Update-PDMD-Progress-3.pdf (accessed 30 July 2011).
- Hamer, D. J. 2011. Update #8—March 2011: Gearing up for sea trials. Non-lethal options for mitigating catch depredation by toothed whales from pelagic longlines. pp. 39–44.

- Available at http://www.marinemammals.gov.au/__data/assets/pdf_file/0004/36976/8.-Update-Gearing-up-for-sea-trials-3.pdf (accessed 30 July 2011).
- Hamer, D. J., T. M. Ward and R. McGarvey. 2008. Measurement, management and mitigation of operational interactions between the South Australian sardine fishery and short-beaked common dolphins (*Delphinus delphis*). *Biological Conservation* 141:2865–2878.
- Hamer, D. J., A. Ivey and T. M. Ward. 2009. Operational interactions of the South Australian sardine fishery with the common dolphin: November 2004 to March 2009. Primary Industry and Resources South Australia. SARDI publication no. F2007/001098-2, Report Series no. 354, Government of South Australia, Adelaide, Australia. 32 pp.
- Hamer, D. J., T. M. Ward, P. D. Shaughnessy and S. R. Clark. 2011. Assessing the effectiveness of the Great Australian Bight Marine Park in protecting the endangered Australian sea lion (*Neophoca cinerea*) from bycatch mortality in shark gill nets. *Endangered Species Research* 14:203–216.
- Hernandez-Milian, G., S. Goetz, C. Varela-Dopico, *et al.* 2008. Results of a short study of interactions of cetaceans and longline fisheries in Atlantic waters: Environmental correlates of catches and depredation events. *Hydrobiologia* 612:251–268.
- Hucke-Gaete, R., C. A. Moreno and C. J. Arata. 2004. Operational interactions of sperm whales and killer whales with the Patagonian toothfish industrial fishery off southern Chile. *CCAMLR Science* 11:127–140.
- Indian Ocean Tuna Commission. 2007. Workshop on the depredation in the tuna long-line fisheries in the Indian Ocean. Report, Indian Ocean Tuna Commission (IOTC), Victoria, Seychelles, 9–10 July 2007 (unpublished). 50 pp. Available at <http://www.iotc.org/files/proceedings/2007/sc/IOTC-2007-SC-INF01.pdf> (accessed 30 July 2011).
- Indian Ocean Tuna Commission. 2010. Kobe II bycatch workshop background paper. Marine mammals. IOTC-2010-WPEB-Inf13. 11 pp. Available at <http://www.iotc.org/files/proceedings/2010/wpeb/IOTC-2010-WPEB-Inf13.pdf> (accessed 30 July 2011).
- Iwashita, M., M. Iuone and Y. Iwasaki. 1963. On the distribution of *Orchinus* in the Northern and Southern Pacific equatorial waters as observed from reports on *Orchinus* predation. *Bulletin of the Fisheries Research Laboratory, Tokai University* 1:24–30.
- Jefferson, T. A. 1994. Marine mammals of the world. Report, United Nations (UN) Food and Agriculture Organisation (FAO), Rome, Italy. 320 pp.
- Jefferson, T. A., and B. E. Curry. 1996. Acoustic methods of reducing or eliminating marine mammal-fishery interactions: Do they work? *Ocean and Coastal Management* 31:41–70.
- Johnson, C. S., M. W. McManus and B. L. Scronce. 1982. Study of fish bite on the AN/BQR-15 towed array. Report, United States Naval Ocean Systems Center, NOSC-TR-867. 23 pp.
- Johnston, D. W., and T. H. Woodley. 1998. A survey of acoustic harassment device (AHD) use in the Bay of Fundy, NB, Canada. *Aquatic Mammals* 24:51–61.
- Kaschner, K. 2004. Modelling and mapping resource overlap between marine mammals and fisheries on a global level. Ph.D. thesis, University of British Columbia, Vancouver, Canada. 240 pp. Available at <http://www2.fisheries.com/archive/grad/abstracts/kaschnerthesis.pdf> (accessed 30 July 2011).
- Kastelein, R. A., N. Jennings, W. C. Verboom, D. De Haan and N. M. Schooneman. 2006. Differences in the response of a striped dolphin (*Stenella coeruleoalba*) and a harbour porpoise (*Phocoena phocoena*) to an acoustic alarm. *Marine Environmental Research* 61:363–378.
- Klein, C. J., A. Chan, L. Kircher, *et al.* 2008. Striking a balance between biodiversity and socioeconomic viability in the design of marine protected areas. *Conservation Biology* 22:691–700.
- Kobe II. 2010. Kobe II workshop on bycatch. Report of the international workshop on tuna RFMO management of issues relating to bycatch. Doc. No. T-RFMO2_W3_5rev1_ENG. 23–25 June 2010. 19 pp. Available at <http://www.wcpfc.int/doc/eb-ip-05/report-international-workshop-tuna-rfmo-management-issues-relating-bycatch> (accessed 30 July 2011).

- Kock, K.-H., M. G. Purves and G. Duhamel. 2006. Interactions between cetacean and fisheries in the Southern Ocean. *Polar Biology* 29:379–388.
- Kraus, S. D., A. J. Read, A. Solow, K. Baldwin, T. Spradlin, E. Anderson and J. Williamson. 1997. Acoustic alarms reduce porpoise mortality. *Nature* 388:525.
- Krutzen, M., W. B. Sherwin, P. Berggren and N. Gales. 2004. Population structure in an inshore cetacean revealed by microsatellite and mtDNA analysis: Bottlenose dolphins (*Tursiops* sp.) in Shark Bay, Western Australia. *Marine Mammal Science* 20:28–47.
- Lauriano, G., L. Caramanna, M. Scarno and F. Andaloro. 2009. An overview of dolphin depredation in Italian artisanal fisheries. *Journal of the Marine Biological Association of the United Kingdom* 89:921–929.
- Leatherwood, S., R. R. Reeves and L. Foster. 1983. *The Sierra Club handbook of dolphins and whales*. Sierra Club Books, San Francisco, CA.
- Le Quesne, W. J. F. 2009. Are flawed MPAs any good or just a new way of making old mistakes? *Journal of Marine Science* 66:132–136.
- Lien, J., C. Hood, D. Pittman, et al. 1995. Sensory systems of aquatic mammals. Pages 349–364 in R. A. Kastelein, J. A. Thomas and P. E. Nachtigall, eds. *Field tests of acoustic devices on groundfish gillnets: Assessment of effectiveness in reducing harbour porpoise bycatch*. De Spil, Woerden, The Netherlands.
- Lukoschek, V., N. Funahashi, S. Lavery, M. L. Dalebout, F. Cipriano and C. S. Baker. 2009. High proportion of minke whales sold on Japanese markets is due to illegal, unreported and unregulated exploitation. *Animal Conservation* 12:385–395.
- Mangel, J. C., J. Alfaro-Shigueto, K. Van Waerbeek, C. Caceres, S. Bearhop, M. J. Witt and B. J. Godley. 2010. Small cetacean captures in Peruvian artisanal fisheries: High despite protective legislation. *Biological Conservation* 143:136–143.
- Matkin, C. O. 1986. Killer whale interactions with the sablefish longline fishery in Prince William Sound, Alaska, 1985, with comments on the Bering Sea. Report, National Marine Mammal Laboratory, National Marine Fishery Service (NMFS), WA98115-0700, 40-HANF-6-0068. 10 pp.
- McPherson, G. R., C. I. Clague, C. R. McPherson, et al. 2008. Reduction of interactions by toothed whales with fishing gear. Phase 1: Development and assessment of depredation mitigation devices around longlines. Report, Fisheries Research and Development Corporation (FRDC), 2003/016. 216 pp. Available at http://www.nmfs.noaa.gov/pr/interactions/fkwtrt/meeting2/mcpherson_et_al_2003.pdf (accessed 30 July 2011).
- Miller, D. L. 2007. *Reproductive biology and phylogeny of cetacea: Whales, porpoises and dolphins*. Science Publishers, Enfield, NH.
- Mitchell, E. D. 1975. Porpoise, dolphin and small whale fisheries of the world: Status and problems. Report, International Union for the Conservation of Nature (IUCN). 129 pp.
- Mooney, T. A., A. F. Pacini and P. E. Nachtigall. 2009. False killer whale (*Pseudorca crassidens*) echolocation and acoustic disruption: Implications for longline bycatch and depredation. *Canadian Journal of Zoology* 87:726–733.
- Moore, J. E., T. M. Cox, R. L. Lewison, et al. 2010. An interview-based approach to assess marine mammal and turtle captures in artisanal fisheries. *Biological Conservation* 143:795–805.
- Moreno, C. A., R. Castro, L. J. Mujica and P. Reyes. 2008. Significant conservation benefits obtained from the use of a new fishing gear in the Chilean Patagonian toothfish fishery. *CCAMLR Science* 15:79–91.
- Morton, A. B., and H. K. Symonds. 2002. Displacement of *Orcinus orca* (L.) by high amplitude sound in British Columbia, Canada. *Journal of Marine Science* 59:71–80.
- Nielsen, B. K., and B. Møhl. 2006. Hull-mounted hydrophones for passive acoustic detection and tracking of sperm whales (*Physeter macrocephalus*). *Applied Acoustics* 67:1175–1186.
- Nishida, T., and M. Tani. 2001. Summary of the depredation surveys for the tuna longline in the Indian and the Pacific Ocean based on the Japanese investigation cruises (1954, 1958 and 1966–81). Working Paper on Tropical Tunas, Indian Ocean Tuna Commission (IOTC). 17 pp.

- Northridge, S. P. 1984. World review of interactions between marine mammals and fisheries. Report #251, United Nations (UN) Food and Agriculture Organization (FAO), Rome, Italy. 190 pp.
- Northridge, S. P. 1991. An updated world review of interactions between marine mammals and fisheries. Report # 251, supplement 1, United Nations (UN) Food and Agriculture Organization (FAO), Rome, Italy. 58 pp.
- Northridge, S. P., and R. J. Hofman. 1999. Marine mammal interactions with fisheries. Pages 99–119 in J. R. Twiss and R. R. Reeves, eds. Conservation and management of marine mammals. Melbourne University Press, Melbourne, Australia.
- Nowacek, D. P., L. H. Thorne, D. W. Johnston and P. L. Tyack. 2007. Response of cetaceans to anthropogenic noise. *Mammal Review* 37:81–115.
- Perves, M. G., D. J. Agnew, E. Baluerias and C. J. Moreno. 2004. Killer whale (*Orcinus orca*) and sperm whale (*Physeter macrocephalus*) interactions with longline vessels in the Patagonian toothfish fishery at South Georgia, South Atlantic. *CCAMLR Science* 11:111–126.
- Pilot, M., M. E. Dalheim and A. R. Heolzel. 2010. Social cohesion among kin, gene flow without dispersal and the evolution of population genetic structure in the killer whale (*Orcinus orca*). *Journal of Evolutionary Biology* 23:20–31.
- Pimper, L. E., C. S. Baker, R. N. P. Goodall, C. Olavarria and M. I. Remis. 2010. Mitochondrial DNA variation and population structure of Commerson's dolphins (*Cephalorhynchus commersonii*) in their southernmost distribution. *Conservation Genetics* 11:2157–2168.
- Price, G. R. 1981. Implications of a critical level in the ear for assessment of noise hazard at high intensities. *Journal of the Acoustical Society of America* 69:171–177.
- Ramos-Cartelle, A., and J. Mejuto. 2008. Interaction of the false killer whale (*Pseudorca crassidens*) and the depredation on the swordfish catches of the Spanish surface longline fleet in the Atlantic, Indian and Pacific Oceans. Report, International Commission for the Conservation of Atlantic Tunas (ICCAT), Collective Volume of Scientific Papers (SCRS/2007/025), 62(6):1721–1783. Available at http://www.iccat.es/Documents/CVSP/CV062_2008/no_6/CV062061721.pdf (accessed 30 July 2011).
- Read, A. J. 2005. Bycatch and depredation. Pages 5–17 in J. E. Reynolds, W. F. Perrin, R. R. Reeves, S. Montgomery and T. J. Ragen, eds. Marine mammal research: Conservation beyond crisis. Johns Hopkins University Press, Baltimore, MD.
- Read, A. J., P. Drinker and S. Northridge. 2006. Bycatch of marine mammals in US and global fisheries. *Biological Conservation* 20:163–169.
- Reeves, R. R., S. Leatherwood and W. F. Perrin. 1994. Dolphins, porpoises and whales. 1994–1998 Action plan for the conservation of cetaceans. Report, IUCN/SSC Cetacean Specialist Group. International Union for the Conservation of Nature, Gland. 91 pp. Available at <http://data.iucn.org/dbtw-wpd/edocs/1994-014.pdf> (accessed 30 July 2011).
- Reeves, R. R., S. Leatherwood and R. W. Baird. 2009. Evidence of possible decline since 1989 in false killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands. *Pacific Science* 63:253–261.
- Roche, C., C. Guinet, N. Gasco and G. Duhamel. 2007. Marine mammals and demersal longline fishery interactions in the Crozet and Kerguelen exclusive economic zones: An assessment of depredation levels. *CCAMLR Science* 14:67–82.
- Secchi, E. R., and T. J. Vaske. 1998. Killer whale (*Orcinus orca*) sightings and depredation on tuna and swordfish longline catches in southern Brazil. *Aquatic Mammals* 24:117–122.
- Secchi, E. R., J. Y. Wang, L. Dalla Rosa, S.-C. Yang and R. R. Reeves. 2005. Global review of interactions between cetaceans and longline fisheries: Preliminary data. Report, International Whaling Commission (IWC), SC/57/SC3. 8 pp.
- Shaughnessy, P. D., R. Kirkwood, M. Cawthorn, C. Kemper and D. Pemberton. 2003. Pinnipeds, cetaceans and fisheries in Australia: A review of operational interactions. Pages 136–152 in N. Gales, M. Hindell and R. Kirkwood, eds. Marine mammals: Fisheries, tourism and management issues. CSIRO Publishing, Melbourne, Australia.
- Shaughnessy, P. D., S. D. Goldsworthy, D. J. Hamer, B. Page and R. R. McIntosh. 2011. Australian sea lions *Neophoca cinerea* at colonies in South Australia: Distribution, abundance and trends, 2004 to 2008. *Endangered Species Research* 13:87–98.

- Sigler, M. F., C. R. Lunsford, J. M. Straley and J. B. Liddle. 2008. Sperm whale depredation of sablefish longline gear in the northeast Pacific Ocean. *Marine Mammal Science* 24: 16–27.
- Sivasubramaniam, K. 1964. Predation of tuna longline catches in the Indian Ocean, by killer whales and sharks. *Bulletin of the Fisheries Research Station* 17:221–236.
- South Pacific Regional Environment Program. 2002. Plan of action and priorities for research to reduce depredation on longlines by cetaceans. Pages 26–34 *in* M. Donoghue, R. R. Reeves and G. Stone, eds. Report of the workshop: Cetacean interactions with commercial longline fisheries in the South Pacific region: Approaches to mitigation. Apia, Samoa, 11–15 November 2002. New England Aquarium Press, Boston, MA. 44 pp. Available at <http://www.nmfs.noaa.gov/pr/pdfs/interactions/samoa2002.pdf> (accessed 30 July 2011).
- Straley, J., T. O'Connell, G. Beam, S. Mesnick, A. Allen and E. Mitchell. 2002. Sperm whale depredation in the demersal longline fishery for sablefish in the Gulf of Alaska: Research needs and approaches to mitigation. Pages 7–8 *in* M. Donoghue, R. R. Reeves and G. Stone, eds. Report of the workshop: Cetacean interactions with commercial longline fisheries in the South Pacific region: Approaches to mitigation. Apia, Samoa, 11–15 November 2002. New England Aquarium Press, Boston, MA. 44 pp. Available at <http://www.nmfs.noaa.gov/pr/pdfs/interactions/samoa2002.pdf> (accessed 30 July 2011).
- Thode, A., J. Straley, C. O. Tiemann, K. Folkert and V. O'Connell. 2007. Observations of potential acoustic cues that attract sperm whales to longline fishing in the Gulf of Alaska. *Journal of the Acoustical Society of America* 122:1265–1277.
- Tilzey, R., S. D. Goldsworthy, M. Cawthorn, *et al.* 2006. Assessment of seal-fishery interactions in the winter blue grenadier fishery off west Tasmania and the development of fishing practices and seal exclusion devices to mitigate seal bycatch by factory trawlers. Report, Fisheries Research and Development Corporation (FRDC), Project no. 2001/008. Bureau of Rural Sciences (BRS). 69 pp. Available at http://adl.brs.gov.au/brsShop/data/frdc_final_reportv2.pdf (accessed 30 July 2011).
- Tixier, P., N. Gasco, G. Duhamel and C. Guinet. 2009. Interactions of Patagonian toothfish fisheries with killer whales and sperm whales in Crozet Exclusive Economic Zone: An assessment of depredation levels and insights on possible mitigation solutions. Report, Secretariat of the Convention for the Conservation of Antarctic Marine Living Resources (CCAMLR), #WG-IMAF-09/12 (unpublished). 30 pp. Available at http://www.ccamlr.org/ccamlr_science/Vol-17-2010/10Tixier-et-al.pdf (accessed 30 July 2011).
- Trippel, E. A., M. B. Strong, J. M. Terhune and J. D. Conway. 1999. Mitigation of harbour porpoise (*Phocoena phocoena*) bycatch in the gillnet fishery in the lower Bay of Fundy. *Canadian Journal of Fisheries and Aquatic Sciences* 56:113–123.
- United Nations. 2009. World population prospects: The 2008 revision. Population Newsletter No. 87, Population Division, United Nations (UN), New York, NY. 20 pp.
- Visser, I. N. 2000. Killer whale (*Orcinus orca*) interactions with longline fisheries in New Zealand waters. *Aquatic Mammals* 26:241–252.
- Wade, P. R. 2002. Population dynamics. Pages 974–979 *in* W. F. Perrin, B. Würsig and J. G. M. Thewissen, eds. *Encyclopedia of marine mammals*. Academic Press, San Diego, CA.
- Ward, P., and S. Hindmarsh. 2007. An overview of historical changes in the fishing gear and practices of pelagic longliners, with particular reference to Japan's Pacific fleet. *Reviews in Fish Biology and Fisheries* 17:501–516.
- Waring, G. T., E. Josephson, C. P. Fairfield and K. M. Foley. 2006. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments—2005. Report, National Oceanographic and Atmospheric Administration (NOAA), NMFS-NE-194, Washington, DC. 346 pp. Available at <http://www.nefsc.noaa.gov/publications/tm/tm213/tm213.pdf> (accessed 30 July 2011).

- Watson J. W., and D. W. Kersletter. 2006. Pelagic longline fishing gear: A brief history and review of research efforts to improve selectivity. *Marine Technology Society Journal* 40:6–11.
- Watson, L. 1981. *Sea guide to the whales of the world*. Dutton Press, London, U.K.
- Williams, A. J., S. L. Petersen, M. Goren and B. P. Watkins. 2007. Sightings of killer whales *Orcinus orca* from longline vessels in South African waters, and consideration of the regional conservation status. *African Journal of Marine Science* 31:81–86.
- Yamaguchi, Y. 1989. Tuna long-line fishing I: Historical aspects. *Marine and Freshwater Behaviour and Physiology* 15:1–11.
- Yano, K., and M. E. Dahlheim. 1995. Killer whale, *Orcinus orca*, depredation of longline catches of bottom fish in the southeastern Bering Sea and adjacent waters. *Fishery Bulletin* 93:355–372.
- Zollett, E. A., and A. J. Read. 2006. Depredation of catch by bottlenose dolphins (*Tursiops truncatus*) in the Florida king mackerel (*Scomberomorus cavalla*) troll fishery. *Fishery Bulletin* 104:343–349.

Received: 3 May 2011
Accepted: 10 August 2011